




# BIODAVERSITY











#### **Seriously?**

#### In the US

40% believe the earth is only 10,000 years old

40% reject evolution

50% deny anthropogenic climate change



### Psychological link found between climate change denial and belief in free market economics

AMERICAN PSYCHOLOGICAL ASSOCIATION — May 4, 2016






UPDATED TO INCLUDE THE LATEST GUIDELINES AND RESEARCH

## The loo,000 copies sold! Paleo Loopies sold! Copies sold!


Lose Weight and Get Healthy
by Eating the Foods
You Were Designed to Eat


Loren Cordain, Ph.D.

Author of The Paleo Diet Cookbook



#### Colonic hydrotherapy





#### FLAT EARTH








Interstellar lizards in people suits dominate the global ruling elite.

Climate change?
Its all rubbish
Fake news.



Of 33,700 authors of peer-reviewed

#### CLIMATE CHANGE



#### 220.000 peer reviewed papers on climate change

- 1/2 reporting on research undertaken
- Team effort of 3 full time equivalents
- Average study time of 2 years


GGU,UUU DEKSUN VEARS OF RESEARCH

#### Human health

**Heart disease** –Kills one Australian every 12 minutes

**Cancer** – 47,753 died if cancer in 2017


**Diabetes** - One person every five minutes diagnosed




#### W.T.F

Scientific research shows an unequivocal link between these diseases and eating animal products.






#### 80% of Americans don't trust science











In defence of science



























Inorganic chemistry, organic chemistry, calculus, statistics, biometrics physics, animal biology, plant biology, plant physiology, computer science, agronomy, entomology, microbiology, genetics, plant pathology, geology, soil science, economics, agricultural economics, plant nutrition, science communication, pasture science, agricultural ecology, natural resource management, agricultural engineering.

#### International Center for Agricultural Research in the Dry Areas













**ICARDA's mission** is to improve the welfare of poor people and alleviate poverty through research and training in dry areas of the developing world to increase the production and nutritional quality of food, while preserving and enhancing the natural resource base.





















**Figure 5.1**: Underground seed softening profiles for 6 accessions (ICARDA accession numbers shown in legend box), which are examples of the variation both in half-lives and the final level of hardseededness of the collection as a whole. The curves represent models of the form (y=A+C/(1+exp[-B(t-M)])) fitted to the seed softening data set generated for each accession. Table 5.4 shows the parameters for each curve with their respective standard errors.

### **Genetic Resource Section (ICARDA-GRS)**

Mission:

collect

conserve

characterize

evaluate


document

distribute

Genetic diversity of wheat, barely, food legumes, pasture and forage species from diverse origins for use in research and plant breeding

















































## Enormous intra-specific diversity





# 80% of the genetic diversity of major crop plants were lost in the last century.

# Examples of Genetic erosion in crop varieties

Стор	Country	Number of varieties in use last century	Number of varieties in use now
Rice	Philippines	1000s	2 (98% of land)
Rice	China	8000 (1949)	50 (1970s)
Rice	Sri Lanka	2000 (1959)	5 (most land)
Wheat	China	10,000 (1949)	1000 (1970s)
Cabbage	USA		95% lost
Field corn	USA		91% lost
Pea	USA		86% lost




































#### Geographic coverage of the ICARDA genetic resource collection





### **Doco Trailer**





## Urban populations eat 8.4 million tons of food per day



## By 2050 global demand for food will increase by 60 - 90%















#### Resources flow

#### N,P,K 200 million tonnes






















6800 trillion litres

### How secure is industrial agriculture?



#### **Inter-connected negative factors**

Climate change

**Population growth** 

Energy crisis/biofuels

Land degradation

Low genetic diversity

**Looming water crisis** 


New pests and diseases

Under-investment in agricultural research

#### Climate change


Likelihood (in percent) that the summer average temperature in 2090 will exceed the highest summer temperature ever observed (1900-2006).

#### Summers in 2080-2100 Warmer than Warmest on Record




Source: Battisti D. S. and Rosamond L. Naylor R. L. 2008. Historical Warnings of Future Food Insecurity with Unprecedented Seasonal Heat. in publication.

#### India



Source: Data from Naylor R., Falcon W. and Fowler C. 2007. The Conservation of Global Crop Genetic Resources In the Face of Climate Change: Summary Statement from a Bellagio Meeting. (Taken from D. Battisti.)






### Water


Use 3 times more water than a generation ago


By 2025 1.8 billion people will live with water shortage

2/3 world in water stress




#### Areas of physical and economic water scarcity





"There is a water crisis today. But the crisis is not about having too little water to satisfy our needs. It is a crisis of managing water so badly that billions of people - and the environment - suffer badly." World Water Vision Report



1 KG potato = 100 liters



1 kg beef = 13,000 liters











#### A paradigm shift



# Innovative science is desperately required











Australian government funding for research has dropped to the lowest it had ever been at just 0.4% of GDP.










### Plant Breeding



- Multiple disease resistance
- Multiple insect resistance
- High water use efficiency
- High nutrient use efficiency
- Drought tolerance
- Abiotic stress tolerance
- Heat tolerance
- Higher yield


# But where do you get the genes?












### Core collections

- Good to explore the general diversity in a collection
- Good to undertake comparative molecular studies
- Fun and interesting to do
- Generate publications 1000s
- But!! how useful are they to actually select subsets of material containing specific traits?
- · Rarely if ever requested by user

#### A matter of LUCK



Depending on the set size and the frequency of the sought after gene getting what you want was most often a question of luck

How do I make a rational decision




How do we find what we want, in a small subset?

How do I take the guesswork out of genebank selections



Faster solutions for crops of tomorrow

# Focused identification of Germplasm Strategy





### FIGS premise

## Emergence of adaptive traits are influenced by environmental factors


































RESEARCH ARTICLE

#### Investigating the case of human nose shape and climate adaptation


Arslan A. Zaidi , Brooke C. Mattern, Peter Claes, Brian McEcoy, Cris Hughes, Mark D. Shriver

Published: March 16, 2017 • https://doi.org/10.1371/journal.pgen.1006616

### FIGS: In a nutshell

Look to environments that are likely to have imposed a selection pressure for the required trait.






Long-term daily averages – GIS

Temp at 2 meter max temp at 2 meter min temp at 2 meter **Precipitation Absolute humidity Relative humidity** Photosynthesis active radiation Wind at east-west direction Wind at north-south direction Vapor pressure deficit

Soil data 109,500 global surfaces generated



Over 200TB of data


### FIGS pathways

#### **User defined trait**






#### **Classify collection sites**



Statistical models




#### Filter collection sites



# Screening for salinity tolerance in bread wheat landraces

VIR,ICARDA,AWCC collection

FIGS subset chosen by mapping collection sites over salinity probabilities



Bread wheat landrace collection sites

# Hydroponic system for salt-tolerance evaluation at VIR, Russia



Method after R.Munns *et al.* 2003. Na and K content measured in 3<sup>rd</sup> leaf after salinity treatment imposed for 10 days






### **Preliminary results**

Set	Leaf Na content range in accessions  mg/g dry weight (min – max)	Average leaf Na content  mg/g dry weight	Number of accessions with Na content below 3 mg/g dry weight  accessions %		Total accessions evaluated
salt	1.16 – 22.46	5.42	67	21	320
core	1.57 – 51.26	10.63	10	3	319





### Sunn pest set selection



Starting with a set of over 16,000 accessions from VIR, ICARDA and AWCC

Selected material collected between 30-45 latitude, 35-80 longitude

Excluded CHN, PAK, IND as only recent reports of Sunn pest in these countries – retained 6,328 accessions

Excluded particularly dry environments - rainfall below 280mm/year, low Aridity Index

Excluded sites where the winters temps fall below -10 degrees – retained 534 accessions

### Sunn pest set selection









# Using FIGS to search for indicators of drought tolerance in Faba bean

- ICARDA hold 9545 faba bean accessions
- Selected material based on measures of moisture availability at collection sites




### Results: clear differences between sets






#### Onset of the temperature & -moisture-limited growing period (1)





#### Applying FIGS within growing season






### Multiple trait state data + Collection site daily data




Non-linear model




Support vector machine, Random forest, Neural networks

### Model attempts to discriminate between trait states



Model parameters iteratively refined


### Trained model used to predictively classify sites



#### **Predictive modelling**



Areas where resistance is likely to occur (dark red)



Russian wheat aphid Sunn Pest Pod borer Powdery mildew – wheat and barley Barley yellow dwarf virus Spot blotch Net blotch Drought tolerance Frost tolerance Beet western yellows virus Chickpea chlorotic stunt virus Alfalfa mosaic virus Been yellow mosaic virus





# Need a paradigm shift in what we eat.















### Input Use Efficiency

- 1. less land and water is required
- 2. Greenhouse gas emissions are lower
- 3. High feed/substrate conversion efficiencies
- 4. Can transform low-value organic by-products into high-quality food or feed
- 5. Climate change resilient
- 6. A lot of medicinal values
- 7. Low energy input







### A NUTRITIONAL SUPERSTAR



3900%

more beta-carotene than carrots

2300%

more iron than spinach





300%

more calcium than fresh milk.

280%

more anti-oxidants than blueberries.





200%

more protein than meat

#### JUST 3 GMS OF SPIRULINA

has more antioxidants and anti-inflammatory properties than **5** servings of vegetables.



## Most nutrient dense food on the planet

### Here is what leading science and health organizations say about spirulina:

"1 lb of spirulina is the equivalent to 1000 lb of vegetables"

"Spirulina is the future of nutrition"

"The most complete balanced and nutritious food on earth"







### A medical marvel

- Strong anti-cancer activity
- Anti-allergy activity.
- Helps in the regulation of cholesterol.
- Helps combat liver damage.
- Help regulate blood pressure.
- Alleviates anaemia.
- Boost immune system.

- Helps fight infections.
- Detoxifies the body of heavy metals and toxins.
- Protects the brain from degenerative diseases.
- Improves mental acuity.
- Improves muscle strength and endurance.
- Helps Manage HIV/AIDS.
- Used to treat radiation sickness.

### Spirulina – high input use efficiency



Yield (kg)/unit of input




















# MUITO OBRIGADO!

